

Citation for published version:

Rommers, N, Teunissen, JW, Cumming, S, Pion, J, Rössler, R, D'Hondt, E, Lenoir, M, Malina, RM & Savelsbergh, G 2020, 'Accuracy of maturity prediction equations in individual elite football players', *Annals of Human Biology*, vol. 47, no. 4, pp. 409-416. <https://doi.org/10.1080/03014460.2020.1783360>

DOI:

[10.1080/03014460.2020.1783360](https://doi.org/10.1080/03014460.2020.1783360)

Publication date:

2020

Document Version

Peer reviewed version

[Link to publication](https://doi.org/10.1080/03014460.2020.1783360)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Accuracy of maturity prediction equations in individual elite football players

Jan Willem Teunissen^{1,2*}, <https://orcid.org/0000-0002-8254-7020>

Nikki Rommers^{2,3,4*}, <https://orcid.org/0000-0003-0311-5009>

Johan Pion^{1,2}, <https://orcid.org/0000-0002-2633-9120>

Sean P. Cumming⁵, <https://orcid.org/0000-0003-1705-9642>

Roland Rössler⁶, <https://orcid.org/0000-0002-6763-0694>

Eva D'Hondt³, <https://orcid.org/0000-0001-5646-2261>

Matthieu Lenoir², <https://orcid.org/0000-0003-3906-1137>

Robert M. Malina⁷ <https://orcid.org/0000-0003-4049-2620>

Geert J.P. Savelsbergh⁸ <https://orcid.org/0000-0001-5795-2828>

*First authorship in equal contribution

Affiliations:

1. Department of Sports and Exercise Studies, HAN University of Applied Sciences, Nijmegen, The Netherlands;

2. Department of Movement and Sports Sciences, Ghent University, Ghent, Belgium;

3. Department of Movement and Sports Sciences, Vrije Universiteit Brussel, Brussels, Belgium;

4. Research Foundation Flanders (FWO), Brussels, Belgium

5. Department of Health, University of Bath, Bath, England

6. Amsterdam Collaboration on Health and Safety in Sports, Amsterdam UMC, Vrije Universiteit Amsterdam, Department of Public and Occupational Health, Amsterdam Movement Sciences, Amsterdam, The Netherlands

7. Department of Kinesiology and Health Education, University of Texas, Austin, Texas, USA

8. Faculty of Human Movement Science, VU University, Amsterdam, The Netherlands

Accuracy of maturity prediction equations in individual elite football players

Abstract

Background: Equations predicting age at peak height velocity (APHV) are often used to assess somatic maturity and to adjust training load accordingly. However, no information is available on the intra-individual accuracy of APHV-estimations over time.

Aim: Purpose of this study is to assess the accuracy of predication equations for the estimation of APHV in individual elite youth soccer players.

Subjects and methods: Anthropometric measurements were conducted at least every three months in 17 adolescent elite football players (11.9 ± 0.8 years) from seasons 2008-2009 to 2011-2012. APHV was estimated at each measurement point by four predominant prediction equations. Predicted APHV was compared to the player's observed APHV using one-sample-t-tests and equivalence-tests. Longitudinal stability was assessed by comparing the linear coefficient of the deviation to zero.

Results: In none of the players, predicted APHV was equivalent to the observed APHV. A difference with a large effect size (Cohen's $d > 0.8$) was found in 87% of the cases. Furthermore, the prediction was not stable over time in 71% of the cases.

Conclusions: None of the assessed prediction equations is accurate in estimating APHV of individual players nor stable over time, which makes it challenging to construct training programmes by predicted time from APHV.

Keywords;

Growth, maturation, soccer, puberty, adolescence

Introduction

Puberty is an important phase in the development of youth athletes. Neuroendocrine alterations associated with puberty influence the growth in size, the timing of the adolescent growth spurt (Marceau et al. 2011), as well as the improvements in strength, power, speed and aerobic and anaerobic fitness (Goswami et al. 2014; Leyhr et al. 2018). However, the timing of peak gains in body mass, strength and power occurs, on average, after peak height velocity (PHV) while peak gains in aerobic fitness occur coincident with PHV (Beunen and Malina 1988; Philippaerts et al. 2006). Moreover, the timing (when specific events associated with maturation occur) and tempo (rate) of growth and maturation varies among individuals and is largely a result of heritable traits (i.e. genes) (Marceau et al. 2011). Accordingly, the development of physical and physiological characteristics may show a fluctuating, non-linear pattern over time (Malina et al. 2005).

Identification, selection, transfer and development of youth athletes are related to differences in individual biological maturity status among high-level youth athletes (Meylan et al. 2010; Malina et al. 2015). Moreover, puberty coincides with a stage of player development where there is increased emphasis on player selection and de-selection and where the physical demands and intensity of training sessions and competition increase (Tierney et al. 2016). A selection bias towards male football players advanced in maturation emerges from approximately 11 years of age and increases with age.(Johnson et al. 2017) In contrast, late maturing players have been shown disproportionally represented in youth football (Johnson et al. 2017).

A commonly used indicator of maturational timing is predicted age at PHV, based on several anthropometric dimensions (Mirwald et al. 2002; Moore et al. 2015; Fransen et al. 2018). The original prediction equation (2002) estimates the maturity offset, an indicator of the time before

or after the age of PHV, from chronological age (CA), height, weight, sitting height and estimated leg length. This prediction equation has since been modified (Moore 1, includes age and sitting height) and simplified by eliminating sitting height from the equation (Moore 2, includes age and height) (Moore et al. 2015). More recently, the linear prediction equation has been extended to a polynomial prediction equation estimating a maturity ratio (Fransen et al. 2018).

The authors of the original equations report error margins around one year in boys (Mirwald et al. 2002; Moore et al. 2015). In addition, several validation analyses in longitudinal samples spanning from 8 through 18 years report major limitations of the original and modified equations. Predicted ages at PHV increase, on average, with CA at prediction, have a reduced range of variation, and have major limitations with early and late maturing youth defined by observed ages at PHV (Malina and Koziel 2014a, 2014b; Malina et al. 2016; Koziel and Malina 2018). At best, the prediction equation may be useful within a narrow CA band among average maturing boys. Consistent with the validation studies, an increase in the average predicted age at PHV was noted in a sample of elite football players 9 and 15 years of age (Rommers et al. 2019).

The preceding observations question the utility of the maturity offset or age at PHV prediction equations for individuals and have implications in the context of individualizing training prescriptions, identifying a player's potential and assessment of injury risk. Hence, the accuracy of predicted ages at PHV in individual youth football players during the interval of adolescence merits attention. In this context, the aim of this study is to investigate the accuracy and longitudinal stability of predicted ages at PHV in elite youth football player who were measured at least every three months during adolescence.

102 **Subjects and methods**

103 ***Participants***

104 Data were collected from seasons 2008-2009 through 2011-2012 in a professional youth
105 football academy in the Netherlands. Players were selected by the academy based on estimated
106 potential in terms of technical, tactical, social and physical skills. All players in the youth
107 academy were measured on a regular basis. Seventeen players (n=17; Caucasian n = 10, African
108 n = 5, Middle Eastern n = 2) were longitudinally followed over at least two years. To ensure a
109 high temporal follow-up around the adolescent growth spurt, only players with at least 15
110 measurement points over an interval that spanned at least two years around the age of PHV
111 were included in this study. After medical checks all participating players were found healthy
112 and had no known growth disorders.

113 ***Procedures***

114 All measurements in this study were part of the regular programme of the club and supervised
115 by the medical staff. All parents and players signed a contract with the club approving their
116 child would take part in the academy's regular programme including professional training and
117 testing and were informed bi-annually on the progress and assessments of their child's
118 performance and growth status. The study followed the principles of the Declaration of
119 Helsinki.

120 ***Anthropometric Assessment***

121 Body dimensions were measured frequently (range every 1 to 6 months) by trained movement
122 scientists prior to a training session in the controlled environment of the dressing rooms.
123 Following the protocol described in Lohman et al. (1988), height was measured (Seca 213i) to

the nearest 0.1 centimetre. Sitting height was measured (Seca 213i) with the player sitting on a stool of standardized height. Sitting height was subtracted from standing height to estimate leg (sub ischial) length. Weight was measured (Seca 803) to the nearest 0,1 kilogram.

Age at peak height velocity

Age at PHV was predicted using the original equation for boys (2002), the two modified equations (2015) and the maturity ratio (2018) (Table 1). The first three equations predict maturity offset; age at PHV is estimated as CA minus predicted offset. With the maturity ratio protocol, CA was divided by the maturity ratio to estimate age at PHV.

*** Insert Table 1 near here ***

Analyses

Descriptive statistics of the first measurement of each player are presented as means with corresponding standard deviations (SD). Age at PHV for individual players was then estimated with Preece-Baines model I (Preece and Baines 1978). The height records of seventeen players were successfully modelled and were used in the analysis.

The deviation between observed age at PHV and predicted ages at PHV with each of the four prediction equations (predicted age at PHV – observed age at PHV) was calculated at each observation for individual players. The observed and the predicted ages at PHV were then compared in each player using one sample t-tests. Subsequently, tests of equivalence using Cohen's d as an effect size, 90% confidence intervals, and pre-determined upper and lower equivalence bounds of ± 0.25 , were calculated to evaluate if the differences were sufficiently sizeable for practical consideration (Lakens et al. 2018). Effect sizes were interpreted as small

145 when Cohen's d was > 0.2 , as moderate with Cohen's d was > 0.5 and as large with Cohen's d
146 was > 0.8 (Cohen 1988).

147 Linear regression was used to investigate the stability of the deviation over the interval of the
148 observations. Due to the small monthly increase in height, monthly measurements and
149 estimated growth velocities are affected by measurement, diurnal and potentially seasonal
150 variability. Therefore, linear regression was used instead of actual data points. To visualize the
151 stability of deviation over the course of the study, regression lines for the four prediction
152 equations were plotted by years from observed PHV for each individual player. If the deviation
153 of the linear coefficient of the regression line for each prediction within individuals was equal
154 to zero, stability of predicted ages at PHV was accepted. All analyses were performed in R
155 (version 3.5.4), with alpha level of significance set at 0.05.

156 In order to visualise the individual growth patterns of the included players, we fitted cubic
157 splines from the age of the first to the last measurement in Microsoft Excel using the SRS1
158 cubic spline software (SRS1 Software, LLC, Boston, MA, USA) with data interpolated to three-
159 month intervals.

160

Results

Predicted and observed APHV

Anthropometric characteristics at baseline are summarized in Table 2.

*** Insert Table 2 near here ***

Observed ages at PHV based on Preece-Baines model I ranged from 12.55 to 15.18 years with mean of 13.8 ± 0.7 years (Table 3). Average predicted ages at PHV based on the four prediction equations of ranged from 13.2 to 15.5 years (Mirwald), from 13.3 to 15.3 years (Moore 1), from 12.9 to 14.8 years (Moore 2) and from 13.2 to 15.1 years (Fransen). The mean and SD of the predicted ages at PHV with each of the four prediction equations for individual players are summarized in Table 3.

*** Insert Table 3 near here ***

The ranges of predicted ages at PHV with each prediction equation for individual players are presented in Table 3. With the Mirwald equation, none of the players showed a mean age of predicted PHV that was equivalent to and not statistically different from the observed age at PHV. There were no instances in which predicted ages at PHV were equivalent to the observed age at PHV. In 87% of the predictions, the predicted ages at PHV were not equivalent to the observed age at PHV with the effect size showing a large effect. In seven players, two predictions were less than observed age at PHV, while in most players, predicted ages at PHV were higher than observed age at PHV.

Longitudinal stability of the predicted ages at PHV

The stability of the deviation of the predicted ages at PHV from observed age at PHV over time is shown for each prediction equation in four randomly selected players in Figure 1. The regression lines depict the deviation of predicted ages at PHV from observed age at PHV over the interval of observation by years from observed PHV at prediction; a horizontal line indicates stable predictions over time. Table 4 shows the range the deviation for each prediction equation and the linear coefficients of the regression lines for each individual player. None of the four equations has a stable prediction over time in more than 45% of the players. The Mirwald and Fransen predictions have more stable predictions than the simplified Moore equations. Overall, the results indicate that a maximum of three predicted ages at PHV in a single individual show relative stability over CA ranges represented in the sample. For most players predicted ages at PHV with only one or two equations show stability, but stable predicted ages at PHV with a specific equation over time vary within and among individuals.

*** Insert Figure 1 ***

*** Insert Table 4 ***

Discussion

Predicted ages at PHV derived with the four prediction equations in a longitudinal sample of elite youth football players differed significantly from and were not equivalent to observed age at PHV estimated with Preece-Baines model I for individual players. Moreover, predicted ages at PHV were not stable in most players across the chronological age span represented in the sample.

Comparison to other studies

Validation studies of the original prediction equation (Mirwald et al. 2002) in longitudinal samples of Polish (Malina and Koziel 2014a) and American (Malina et al. 2016) boys and of the modified equations (Moore et al. 2015) in the Polish boys (Koziel and Malina 2018) showed, on average, reduced variation in predicted compared to observed ages at PHV, later predicted than observed ages at PHV in early maturing boys and earlier predicted than observed ages at PHV in late maturing boys. Moreover, cross-sectional studies of elite football players have indicated advanced skeletal and sexual maturity status compared to the general population (Malina 2011; Malina et al. 2012). Nevertheless, allowing inter-individual differences in biological maturity status and timing, intra-individual variation in predicted ages at PHV is considerable and relatively few predictions approximated observed age at PHV (Koziel and Malina 2018).

The initial study, on the Mirwald equation in Polish boys showed, on average, a stable deviation between predicted and observed ages at PHV in *average* maturing boys between 13 and 15 years of age (Malina and Koziel 2014a). This was not consistent with observations for 15 of the 17 boys in our sample who had an observed age at PHV that could be classified as average. A possible explanation for the difference is the frequency of measurements in the present study

compared to annual observations the study of Polish boys (Malina & Koziel, (2018). On the other hand, it is possible that predictions in the present study were affected by measurement variability in height, weight and sitting height across observations in addition to seasonal fluctuations in growth in height and weight. Growth in height is also generally more rapid in the spring/summer and slower in the fall/winter, while growth in weight shows the opposite season pattern (Cole 1998). Seasonal variation in growth may affect predictions made across the football season. It has also been suggested that growth in height occurs in mini-spurts followed by intervals of no increase (Lampl and Johnson 1993).

The prediction equation of Fransen et al. (2018) was validated in a mixed-longitudinal sample of elite youth football players, and as such it was expected that the prediction equation would yield more reliable results. This, however, was not the case in the present study.

Strengths and limitations

The strength of this study may be the high frequency of measurements during the interval of the adolescent growth spurt which permitted a closer evaluation of the growth elite football players. On the other hand, the high frequency of measurements is also a limitation from the perspective measurement variability (inter- and intra-observer) in direct (height, sitting height, weight) and derived (estimated leg length) variables, and the relatively close intervals between measurements. As noted earlier, other potential confounding factors are diurnal and seasonal variation in growth. In addition, estimates of growth rate over short intervals have a larger variance (Tanner et al. 1966; Roche and Himes 1980). It should also be acknowledged that the Preece-Baines model I is a mathematical growth model that has an error margin. This model, however indicates a clear estimate of the age at PHV, which is not the case for cubic splines for example, showing several peaks in some individuals (see Figure 2).

Although the majority of players in our sample were of Caucasian origin, we also included players of different ethnicity. The variation in ethnicity is representative for contemporary elite-level youth football teams. This is of relevance as the prediction equations as well as Preece-Baines model I were based on samples of European ancestry, while ethnic variation in the proportion of leg length to stature is well documented (Malina et al. 2004). As such, care is warranted in generalizing the observations, although they were consistent with several validation studies of the maturity offset/predicted age at PHV protocol.

Practical recommendations for training and future directions

Puberty is a critical period of talent development (Lloyd et al. 2014; Malina et al. 2015). However, it is characterized by considerable inter-individual variation in the timing of the growth spurt in body size and also several indicators of fitness – strength, explosive power and aerobic power in males, both athletes and non-athletes (Philippaerts et al. 2006).

Some evidence indicates a peak incidence of injury around the predicted time of PHV (van der Sluis et al. 2015; Read et al. 2017). It is common to decrease the workload and adjust exercises during the interval of PHV and to focus on individualized training plans (Lloyd and Oliver 2013; Lloyd et al. 2016). For optimal management of training load and in order to maximise athlete development during the interval of PHV, the importance of continuous assessment of growth of youth athletes during the pubertal period has been suggested (Lloyd et al. 2014). Given the non-invasiveness, time and cost efficiency, and immediate outcome predicted maturity offset and/or age at PHV is attractively simple and is increasingly, if not uncritically, used to individualise training and competition programmes (Cumming et al. 2017). However, as shown in the present study, the individual accuracy of all four prediction equations for estimating a player's age at PHV is questionable, and use of the prediction equations in this context is not recommended.

266 Growth in height during adolescence varies considerably among individuals. This individuality
267 of somatic growth emphasizes the need to closely monitor growth status in order to establish
268 training goals. In this context, it is recommended that youth players should be measured at
269 three-month intervals in order to establish meaningful changes and to minimize the influence
270 of daily fluctuations and measurement variability (Lloyd et al. 2014). Such measures can be
271 taken in conjunction with estimates of maturity status to provide a more comprehensive picture
272 of growth and maturity status. Monitoring growth velocity is relatively easy to establish in
273 practice and has the advantage that it considers the non-linear character of growth.
274 Nevertheless, attention to potential seasonal variation in growth should not be overlooked.
275 Future research could focus on adapting training goals and modalities relative to estimated
276 velocities of growth in height during the interval of the adolescent spurt and specific stage of
277 pubertal development (pubic hair, genital) in an effort to individualize training. The authors
278 would like to propose the hypothesis that more frequent assessments of growth will show ‘mini-
279 growth-spurts’ (Figure 2), despite the limitations of the reliability of frequent measurements,
280 and will make it possible to adjust training programs (i.e. intensity, volume and training
281 forms/activities) accordingly. If ‘mini-growth-spurts can be confirmed in future studies than we
282 would like to suggest revising the bio-banding concept by constructing the bands around the
283 rate of growth-velocity rather than maturity offset or percentage of adult stature. Moreover, to
284 support the practitioner in the future with a more accurate tool to assess growth spurts during
285 puberty for 1) the design of athlete development programmes and 2) the assessment of injury
286 risk - a combination of the current equations and growth velocity tracking can be more a valid
287 option. The challenges of this concept are: 1) it is not known how accurate individual
288 extrapolations of frequently measured growth data in the past are for the growth curve ahead,
289 and 2) no cut-off growth-velocity-rates are established by which coaches can adjust their

training. Although his concept might be audacious, it might help the coach to more accurately guide individual pathways of athletes during their transition from adolescents to adults.

Conclusion and practical implications

The results of this longitudinal study in elite youth football players suggested that none of the four equations for predicting age at PHV provides an accurate prediction in individuals. The stability of predictions within individuals was also poor. By inference, the utility of the prediction equations has major limitations. Therefore, we do not recommend the use of the prediction equations to prescribe individualized training programmes or to assess injury risk in youth elite level football players. Future studies could focus on the evaluation of the reliability of frequent measurements of growth (growth tracking) in order to capture possible ‘mini-growth-spurts’ and to assess the associated injury risk and optimal training accordingly.

Acknowledgements

The authors would like to thank the following sport professionals and scientists for their contribution to the data collection and data management; Rosanne Briggeman, Vosse de Boode, Rene Wormhoudt and Olav Versloot.

Declaration of interest statement

The authors report no conflict of interest.

References

- Beunen G, Malina RM. 1988. Growth and physical performance relative to the timing of the adolescent spurt. *Exercise and sport sciences reviews*. 16:503-540. Eng.
- Cohen J. 1988. *Statistical Power Analysis for the Behavioral Sciences*, 2nd Edition. Hillsdale, NJ: Lawrence Erlbaum.
- mCole T. 1998. Seasonality of growth. In «*The Cambridge Encyclopedia of Human Growth and Development*». Ed. by SJ Ulijaszek, FE Johnston & MA Preece. Cambridge Univ. Press, Cambridge.
- Cumming SP, Lloyd RS, Oliver JL, Eisenmann JC, Malina RM. 2017. Bio-banding in Sport. *Strength and Conditioning Journal*. 39(2):34-47.
- Fransen J, Bush S, Woodcock S, Novak A, Deprez D, Baxter-Jones ADG, Vaeyens R, Lenoir M. 2018. Improving the Prediction of Maturity From Anthropometric Variables Using a Maturity Ratio. *Pediatric exercise science*. 30(2):296-307. eng.
- Goswami B, Roy AS, Dalui R, Bandyopadhyay A. 2014. Impact of Pubertal Growth on Physical Fitness. *American Journal of Sports Science and Medicine*. 2(5A):34-39.
- Johnson A, Farooq A, Whiteley R. 2017. Skeletal maturation status is more strongly associated with academy selection than birth quarter. *Science and Medicine in Football*. 1(2):157-163.
- Kozziel SM, Malina RM. 2018. Modified Maturity Offset Prediction Equations: Validation in Independent Longitudinal Samples of Boys and Girls. *Sports medicine (Auckland, NZ)*. 48(1):221-236.
- Lakens D, Scheel AM, Isager PM. 2018. Equivalence testing for psychological research: A tutorial. *Advances in Methods and Practices in Psychological Science*. 2515245918770963.
- Lampl M, Johnson ML. 1993. A case study of daily growth during adolescence: a single spurt or changes in the dynamics of saltatory growth? *Annals of human biology*. 20(6):595-603.

332 Leyhr D, Kelava A, Raabe J, Honer O. 2018. Longitudinal motor performance development
 333 in early adolescence and its relationship to adult success: An 8-year prospective study of
 334 highly talented soccer players. PloS one. 13(5):e0196324. eng.

335 Lloyd RS, Cronin JB, Faigenbaum AD, Haff GG, Howard R, Kraemer WJ, Micheli LJ, Myer
 336 GD, Oliver JL. 2016. National Strength and Conditioning Association Position Statement on
 337 Long-Term Athletic Development. Journal of strength and conditioning research. 30(6):1491-
 338 1509. eng.

339 Lloyd RS, Oliver J. 2013. The impact of growth and maturation on physical performance.
 340 Strength and Conditioning for Young Athletes. Routledge; p. 25-40.

341 Lloyd RS, Oliver JL, Faigenbaum AD, Myer GD, De Ste Croix MB. 2014. Chronological age
 342 vs. biological maturation: implications for exercise programming in youth. Journal of strength
 343 and conditioning research. 28(5):1454-1464. eng.

344 Lohman TG, Roche, A.F., Martorell, R. 1988. Anthropometric standardization reference
 345 manual. Champaign, IL: Human Kinetic Books.

346 Malina RM. 2011. Skeletal age and age verification in youth sport. Sports medicine
 347 (Auckland, NZ). 41(11):925-947.

348 Malina RM, Bouchard C, Bar-Or O. 2004. Growth, maturation, and physical activity. Human
 349 kinetics.

350 Malina RM, Choh AC, Czerwinski SA, Chumlea WC. 2016. Validation of Maturity Offset in
 351 the Fels Longitudinal Study. Pediatric exercise science. 28(3):439-455. eng.

352 Malina RM, Coelho ESMJ, Figueiredo AJ, Carling C, Beunen GP. 2012. Interrelationships
 353 among invasive and non-invasive indicators of biological maturation in adolescent male
 354 soccer players. Journal of sports sciences. 30(15):1705-1717. Eng.

355 Malina RM, Cumming SP, Morano PJ, Barron M, Miller SJ. 2005. Maturity status of youth
 356 football players: a noninvasive estimate. *Medicine and science in sports and exercise*.
 357 37(6):1044-1052. Eng.

358 Malina RM, Koziel SM. 2014a. Validation of maturity offset in a longitudinal sample of
 359 Polish boys. *Journal of sports sciences*. 32(5):424-437. eng.

360 Malina RM, Koziel SM. 2014b. Validation of maturity offset in a longitudinal sample of
 361 Polish girls. *Journal of sports sciences*. 32(14):1374-1382. eng.

362 Malina RM, Rogol AD, Cumming SP, Coelho e Silva MJ, Figueiredo AJ. 2015. Biological
 363 maturation of youth athletes: assessment and implications. *British journal of sports medicine*.
 364 49(13):852-859. eng.

365 Marceau K, Ram N, Houts RM, Grimm KJ, Susman EJ. 2011. Individual differences in boys'
 366 and girls' timing and tempo of puberty: modeling development with nonlinear growth models.
 367 *Dev Psychol*. 47(5):1389-1409.

368 Meylan C, Cronin J, Oliver J, Hughes M. 2010. Talent Identification in Soccer: The Role of
 369 Maturity Status on Physical, Physiological and Technical Characteristics. *International*
 370 *Journal of Sports Science & Coaching*. 5(4):571-592.

371 Mirwald RL, Baxter-Jones AD, Bailey DA, Beunen GP. 2002. An assessment of maturity
 372 from anthropometric measurements. *Medicine and science in sports and exercise*. 34(4):689-
 373 694. eng.

374 Moore SA, McKay HA, Macdonald H, Nettlefold L, Baxter-Jones AD, Cameron N, Brasher
 375 PM. 2015. Enhancing a Somatic Maturity Prediction Model. *Medicine and science in sports*
 376 *and exercise*. 47(8):1755-1764. eng.

377 Philippaerts RM, Vaeyens R, Janssens M, Van Renterghem B, Matthys D, Craen R, Bourgois
 378 J, Vrijens J, Beunen G, Malina RM. 2006. The relationship between peak height velocity and
 379 physical performance in youth soccer players. *Journal of sports sciences*. 24(3):221-230. Eng.

380 Preece MA, Baines MJ. 1978. A new family of mathematical models describing the human
 381 growth curve. *Annals of human biology*. 5(1):1-24.
 382 Read PJ, Oliver JL, De Ste Croix MBA, Myer GD, Lloyd RS. 2017. An audit of injuries in six
 383 english professional soccer academies. *Journal of sports sciences*.1-7. eng.
 384 Roche AF, Himes JH. 1980. Incremental growth charts. *Am J Clin Nutr*. 33(9):2041-2052.
 385 Rommers N, Mostaert M, Goossens L, Vaeyens R, Witvrouw E, Lenoir M, D'Hondt E. 2019.
 386 Age and maturity related differences in motor coordination among male elite youth soccer
 387 players. *Journal of sports sciences*. 37(2):196-203. eng.
 388 Tanner JM, Whitehouse R, Takaishi M. 1966. Standards from birth to maturity for height,
 389 weight, height velocity, and weight velocity: British children, 1965. I. *Archives of Disease in*
 390 *Childhood*. 41(219):454.
 391 Tierney PJ, Young A, Clarke ND, Duncan MJ. 2016. Match play demands of 11 versus 11
 392 professional football using Global Positioning System tracking: Variations across common
 393 playing formations. *Hum Mov Sci*. 49:1-8. eng.
 394 van der Sluis A, Elferink-Gemser MT, Brink MS, Visscher C. 2015. Importance of peak
 395 height velocity timing in terms of injuries in talented soccer players. *International journal of*
 396 *sports medicine*. 36(4):327-332. eng.

397

398

399

400 **Table 1.** Overview of the maturity offset/maturity ratio prediction equations for boys

Mirwald	Maturity offset = $-9.236 + (0.0002708 \times \text{leg length} \times \text{sitting height}) - (0.001663 \times \text{age} \times \text{leg length}) + (0.007216 \times \text{age} \times \text{sitting height}) + (0.02292 \times (\text{weight by height ratio}) \times 100)$
Moore 1	Maturity offset = $-8.128741 + (0.0070346 \times (\text{age} \times \text{sitting height}))$
Moore 2	Maturity offset = $-7.999994 + (0.0036124 \times (\text{age} \times \text{height}))$
Fransen	Maturity ratio = $6.986547255416 + (0.115802846632 \times \text{age}) + (0.001450825199 \times \text{age}^2) + (0.004518400406 \times \text{weight}) - (0.000034086447 \times \text{weight}^2) - (0.151951447289 \times \text{height}) + (0.000932836659 \times \text{height}^2) - (0.000001656585 \times \text{height}^3) + (0.032198263733 \times \text{leg length}) - (0.000269025264 \times \text{leg length}^2) - (0.000760897942 \times (\text{height} \times \text{age}))$

401

402

Table 2. Baseline characteristics of the players

	Mean	SD	Range
Age (y)	11.9	0.8	10.9 – 14.1
Height (cm)	149.7	6.2	139.5 – 165.5
Weight (kg)	38.9	5.9	33.0 – 56.0
Sitting Height (cm)	75.8	2.8	70.7 – 82.1
Observed APHV (y)	13.8	0.7	12.6 – 15.2
Number of measurements	19.8	2.3	16 – 25

y: years, cm: centimetre, kg: kilogram, APHV: age at peak height velocity

411

412 **Table 3.** Observed age at PHV (years) compared to predicted ages at PHV (years) with four equations

413

Observed age at PHV	Mirwald		Moore 1		Moore 2		Fransen	
	Range	Cohen's d [90% CI]	Range	Cohen's d [90% CI]	Range	Cohen's d [90% CI]	Range	Cohen's d [90% CI]
12.6	13.3 : 13.7	7.5 [5.2 : 9.5]***	13.5 : 14.0	9.5 [6.7 : 12.2]***	13.2 : 13.3	10.9 [7.6 : 13.9]***	13.1 : 13.6	4.3 [3.0 : 5.5]***
13.0	13.5 : 14.4	3.3 [2.4 : 4.2]***	13.6 : 14.2	4.7 [3.4 : 6.0]***	13.4 : 13.6	9.9 [7.2 : 12.4]***	13.4 : 14.7	2.1 [1.4 : 2.7]***
13.2	13.2 : 13.5	1.5 [0.9 : 2.0]***	13.4 : 14.1	3.8 [2.6 : 4.8]***	13.0 : 13.4	0.5 [0.1 : 0.9]*	13.0 : 13.4	0.1 [-0.3 : 0.5]
13.3	13.4 : 14.0	3.8 [2.8 : 4.8]***	13.5 : 14.2	3.6 [2.7 : 4.6]***	13.1 : 13.7	0.8 [0.4 : 1.2]**	13.4 : 14.3	2.3 [1.6 : 3.0]**
13.4	13.6 : 14.2	3.1 [2.2 : 4.0]***	13.9 : 14.6	4.3 [3.1 : 5.4]***	13.5 : 13.8	3.9 [2.8 : 5.0]***	13.4 : 14.2	2.1 [1.4 : 2.7]***
13.4	13.4 : 13.8	1.9 [1.2 : 2.5]***	13.4 : 13.9	2.2 [1.5 : 2.9]***	13.0 : 13.3	2.0 [1.3 : 2.7]***	13.3 : 14.0	0.7 [0.3 : 1.1]**
13.4	13.8 : 14.1	6.5 [4.7 : 8.3]***	13.4 : 13.9	2.4 [1.7 : 3.1]***	13.3 : 13.5	0.5 [0.1 : 0.8]	13.8 : 14.3	3.7 [2.6 : 4.7]***
13.5	13.6 : 14.2	1.5 [0.9 : 2.1]***	14.0 : 14.7	4.9 [3.4 : 6.4]***	13.7 : 13.8	6.1 [4.2 : 7.8]***	13.4 : 14.2	0.9 [0.4 : 1.4]**
13.6	13.0 : 13.4	3.3 [2.3 : 4.2]***	13.2 : 13.5	4.0 [2.9 : 5.1]***	12.8 : 13.1	9.9 [7.1 : 12.5]***	13.1 : 14.2	0.5 [0.1 : 0.9]*
13.7	13.7 : 14.6	4.2 [3.1 : 5.3]***	13.8 : 14.5	2.7 [1.9 : 3.5]***	13.3 : 14.0	0.9 [0.5 : 1.3]***	13.8 : 14.6	4.4 [3.2 : 5.6]***
14.1	14.1 : 14.5	1.2 [0.7 : 1.8]***	13.9 : 14.2	0.1 [-0.3 : 0.5]	13.6 : 13.8	11.5 [7.8 : 14.8]***	14.1 : 14.6	1.5 [0.9 : 2.1]***
14.1	13.1 : 13.7	3.7 [2.6 : 4.8]***	13.3 : 14.1	0.9 [0.4 : 1.3]**	13.1 : 13.4	6.9 [4.9 : 8.8]***	13.0 : 13.6	4.7 [3.4 : 6.0]***
14.1	13.0 : 13.9	2.7 [1.8 : 3.5]***	14.2 : 14.7	3.0 [2.0 : 4.0]***	13.4 : 13.7	7.5 [5.2 : 9.7]***	13.0 : 13.8	3.5 [2.4 : 4.5]***
14.6	14.6 : 15.4	2.6 [1.7 : 3.4]***	13.8 : 14.3	4.6 [3.2 : 5.9]***	13.7 : 14.0	9.0 [6.3 : 11.4]***	14.7 : 15.7	2.6 [1.7 : 3.4]***
14.6	13.9 : 14.7	0.6 [0.3 : 1.0]**	14.0 : 14.8	0.6 [0.3 : 1.0]**	13.5 : 14.1	4.8 [3.6 : 6.0]***	14.0 : 14.7	1.1 [0.7 : 1.5]***
15.2	14.6 : 15.1	2.3 [1.6 : 2.9]***	14.4 : 15.4	0.4 [0.0 : 0.7]	14.1 : 14.6	4.0 [2.9 : 5.0]***	14.5 : 14.8	5.7 [4.2 : 7.2]***
15.2	15.2 : 15.6	3.2 [2.3 : 4.0]***	15.2 : 15.5	1.0 [0.6 : 1.5]***	14.6 : 14.8	11.4 [8.3 : 14.3]***	15.0 : 15.3	0.8 [0.4 : 1.2]**

428 PHV: peak height velocity, 90% CI: 90% confidence interval, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$

429 **Table 4.** Range of deviation between observed and predicted ages at PHV (years) and slopes of regression lines of predictions over time with each
430 of the four prediction equations.

Observed age at PHV	Mirwald			Moore 1			Moore 2			Fransen		
	Deviation range	Linear coefficient [95% CI]	p- value	Deviation range	Linear coefficient [95% CI]	p- value	Deviation range	Linear coefficient [95% CI]	p- value	Deviation range	Linear coefficient [95% CI]	p- value
12.6	0.73 : 1.13	-0.07 [-0.12 : -0.01]	0.031	0.98 : 1.5	0.13 [0.1 : 0.15]	0.000	0.56 : 0.8	0.05 [0.03 : 0.07]	0.001	0.55 : 1.08	-0.16 [-0.21 : -0.1]	0.000
13.0	0.50 : 1.40	-0.26 [-0.31 : -0.2]	0.000	0.53 : 1.21	0.17 [0.13 : 0.22]	0.000	0.38 : 0.53	-0.01 [-0.03 : 0.01]	0.415	0.33 : 1.63	-0.38 [-0.47 : -0.29]	0.000
13.2	-0.04 : 0.31	-0.01 [-0.07 : 0.04]	0.637	0.2 : 0.86	0.16 [0.1 : 0.22]	0.000	-0.21 : 0.14	0.08 [0.05 : 0.12]	0.000	-0.24 : 0.23	-0.09 [-0.14 : -0.03]	0.004
13.3	0.14 : 0.71	0.03 [-0.02 : 0.08]	0.228	0.2 : 0.86	0.12 [0.1 : 0.15]	0.000	-0.2 : 0.36	0.07 [0.05 : 0.1]	0.000	0.06 : 1.03	0.12 [0.06 : 0.18]	0.001
13.4	0.22 : 0.83	-0.12 [-0.18 : -0.07]	0.000	0.54 : 1.21	0.2 [0.16 : 0.24]	0.000	0.16 : 0.44	0.05 [0.03 : 0.08]	0.001	0.08 : 0.81	-0.16 [-0.22 : -0.1]	0.000
13.4	0.06 : 0.44	-0.1 [-0.13 : -0.06]	0.000	0.01 : 0.52	0.03 [-0.04 : 0.11]	0.374	-0.34 : -0.01	-0.03 [-0.06 : 0.01]	0.154	-0.02 : 0.6	-0.05 [-0.14 : 0.03]	0.213
13.4	0.41 : 0.72	-0.05 [-0.09 : -0.01]	0.019	0 : 0.52	0.14 [0.08 : 0.19]	0.000	-0.16 : 0.07	0.04 [0.02 : 0.07]	0.004	0.34 : 0.88	-0.11 [-0.17 : -0.05]	0.002
13.5	0.11 : 0.75	-0.29 [-0.35 : -0.24]	0.000	0.58 : 1.26	0.25 [0.21 : 0.28]	0.000	0.21 : 0.36	0.01 [-0.02 : 0.04]	0.379	-0.02 : 0.74	-0.33 [-0.39 : -0.28]	0.000
13.6	-0.57 : -0.13	-0.09 [-0.13 : -0.04]	0.002	-0.39 : -0.08	0.05 [0.02 : 0.09]	0.005	-0.75 : -0.51	-0.01 [-0.04 : 0.03]	0.583	-0.47 : 0.58	0.25 [0.17 : 0.33]	0.000
13.7	-0.03 : 0.86	0.08 [0.01 : 0.15]	0.026	0.05 : 0.75	0.15 [0.12 : 0.18]	0.000	-0.42 : 0.24	0.11 [0.08 : 0.14]	0.000	0.09 : 0.9	0.04 [-0.03 : 0.11]	0.224
14.1	-0.01 : 0.39	-0.12 [-0.19 : -0.04]	0.005	-0.18 : 0.15	0.04 [-0.03 : 0.11]	0.241	-0.44 : -0.31	-0.03 [-0.05 : 0]	0.062	0.08 : 0.48	0 [-0.1 : 0.1]	0.998
14.1	-0.93 : -0.33	-0.01 [-0.1 : 0.09]	0.855	-0.7 : 0.06	0.2 [0.12 : 0.28]	0.000	-0.99 : -0.63	0.1 [0.05 : 0.14]	0.000	-1.06 : -0.48	0.01 [-0.08 : 0.1]	0.771
14.1	-1.1 : -0.15	-0.29 [-0.37 : -0.21]	0.000	0.14 : 0.68	0.13 [0.11 : 0.16]	0.000	-0.61 : -0.37	-0.05 [-0.08 : -0.03]	0.001	-1.04 : -0.28	-0.2 [-0.28 : -0.12]	0.000
14.6	0.01 : 0.79	-0.01 [-0.1 : 0.09]	0.879	-0.71 : -0.31	0.09 [0.05 : 0.13]	0.000	-0.83 : -0.51	0.04 [0 : 0.07]	0.031	0.13 : 1.09	-0.03 [-0.15 : 0.08]	0.571
14.6	-0.73 : 0.04	0.01 [-0.05 : 0.07]	0.728	-0.64 : 0.16	0.19 [0.16 : 0.22]	0.000	-1.14 : -0.56	0.1 [0.07 : 0.13]	0.000	-0.62 : 0.02	-0.03 [-0.08 : 0.03]	0.294
15.2	-0.59 : -0.09	0.12 [0.09 : 0.16]	0.000	-0.75 : 0.19	0.28 [0.22 : 0.34]	0.000	-1.12 : -0.53	0.19 [0.16 : 0.22]	0.000	-0.68 : -0.35	0.07 [0.04 : 0.09]	0.000
15.2	0.08 : 0.42	0.02 [-0.04 : 0.07]	0.472	-0.01 : 0.32	0.09 [0.04 : 0.15]	0.002	-0.52 : -0.35	0.05 [0.04 : 0.06]	0.000	-0.15 : 0.11	-0.03 [-0.05 : 0.05]	0.909

431 PHV: peak height velocity, 95% CI: 95% confidence interval

Figure 1. Deviation between observed ages at PHV and predicted age at PHV (years) in four randomly selected players by years from PHV at prediction with each of the four equations.

Black: Mirwald equation, Blue: Moore 1 equation, Red: Moore 2 equation, Grey: Fransen equation




Figure 2. Growth velocity of individual players modelled by cubic splines in four randomly selected players (same players as in figure 1).

